

MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-1963-A



CENTER FOR STOCHASTIC PROCESSES

Department of Statistics University of North Carolina Chapel Hill, North Carolina





A Trivariate Version of 'Levy's Equivalence'

by

Gordon Simons

Technical Report No. 90

Aprilia : 50 dist.

February 1985

OTIC FILE CO

AD-A153

85 41 091

	 ITV	Ĉ.	ARŜI	 CA'	TIÔN.	OF	THIS	PA	GE

				REPORT DOCUME	NTATION PAGE					
1. REPORT	SECURITY C	LASSIFICA	TION		1b. RESTRICTIVE MARKINGS					
20. SECURIT	TY CLASSIFIC	ATION AU	THORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution					
26. DECLAS	SIFICATION/	DOWNGRA	DING SCHED	PULE	unlimited.					
4. PERFORM	ING ORGAN	ZATION RE	PORT NUM	BER(S)	5. MONITORING ORGANIZATION REPORT NUMBER(S)					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) TR No. 90					AFOSR-TR- 85-0295					
6a NAME OF PERFORMING ORGANIZATION University of North				Sb. OFFICE SYMBOL (If applicable)	76. NAME OF MONITORING ORGANIZATION					
Carol:				<u> </u>	Air Force Office of Scientific Research					
6c. ADDRES	S (City, State	end ZIP Cod	Process	es Department	7b. ADDRESS (City, State and ZIP Code)					
Center for Stochastic Processes, Department of Statistics, Chapel Hill NC 27514					Directorate of Mathematical & Information Sciences, Bolling AFB DC 20332-6448					
	F FUNDING/	SPONSORIN	G	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER					
AFOSR				NM	F49620-82-C-0009					
Sc ADDRES	S (City, State	and ZIP Cod	e)		10. SOURCE OF FUNDING NOS.					
					PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT		
	AFB DC				61102F	2304	A5			
	Include Securit VARIATE V			'S EQUIVALENCE'						
	AL AUTHOR	(S)								
134 TYPE C			136. TIME C		14. DATE OF REPORT (Yr., Mo., Day) 15. PAGE COUNT					
Techn:	ical Mentary No	0747101	FROM	<u> </u>	FEB 85		2	2		
10. 307762	MENIAN I N	01211011								
17.	COSATI	CODES		18. SUBJECT TERMS (C	ontinue on reverse if ne	cessary and identi	/y by block numbe	7)		
FIELD	GROUP	SUE	. GR.	Wiener process	Wiener process; Levy's equivalence; Tanaka's formula.					
				<u> </u> 						
19. ABSTRA	CT (Continue	on reverse if	necessary and	d identify by block number	·,		· · · · · · · · · · · · · · · · · · ·			
	It is si	hown tha	t the tr	ivariate stocha	stic processes	: {(MWM.	Θ. t > 0	and		
{(IW	. I. L. T	.1. t >	O} have	the same distril	outions when W		t't' -	200		
proc	t''	t'' — is the m	aximum v	alue attained by	v W over the t	ime interv		ic the		
time	the may	imum val	up ic at	tained l is th	ne local time	of Wat In	21 [U,U], 0	15 the		
time the maximum value is attained, L is the local time of W at level zero and time t,										
and T _t is the last time W is zero in the time interval [0,t]. A straightforward proof, based on "Tanaka's formula", establishes this result by deriving an almost sure version										
			ormara ,	establishes the	is result by d	eriving an	almost sure	e version		
of the equivalence.										
20. DISTRI	BUTION/AVA	ILABILITY	OF ABSTRA	ст	21. ABSTRACT SECURITY CLASSIFICATION					
UNCLASSIF	HED/UNLIMI	TED E SA	ME AS RPT.	□ DTIC USERS □	UNCLASSIFIED					
220. NAME	OF RESPONS	IBLE INDIV	IDUAL		22b. TELEPHONE NUMBER 22c. OFFICE SYMBOL (Include Area Code)			MBOL		
MAJ B	rian W. W	loodruff			(000) 545	5027	NM			

A Trivariate Version of 'Lévy's Equivalence'

by Gordon Simons*
University of North Carolina, Chapel Hill



Summary

It is shown that the trivariate stochastic processes $\{(M_t - W_t, M_t, \Theta_t), t \ge 0\}$ and $\{(|W_t|, L_t, T_t), t \ge 0\}$ have the same distributions when: $W = \{W_t, t \ge 0\}$ is a Wiener process, M_t is the maximum value attained by W over the time interval [0,t], Θ_t is the time the maximum value is attained, L_t is the local time of W at level zero and time t, and T_t is the last time W is zero in the time interval [0,t]. A straightforward proof, based on "Tanaka's formula", establishes this result by deriving an almost sure version of the equivalence.

AMS 1980 subject classification. Primary 60G17 Secondary 60H05

Keywords and phrases. Wiener process, Lévy's equivalence, Tanaka's formula

*This research has been supported by the National Science Foundation Grant No. DMS-8400602.

AIR FORCE OFFICE OF SCIENTIFIC RECEIVED (ADDO)
NOTION OF SAME AND ADDRESS OF SCIENTIFIC RECEIVED (ADDO)
This sapprov
Districts
MATTHEW J. Add.
Chief, Technical Information Division

As described by Knight (1981), "Lévy's equivalence" refers to the equality in distribution of the bivariate stochastic processes $\{(M_{t} - W_{t}, M_{t})\}$ M_t), $t \ge 0$ and $\{(|W_t|, L_t) \ t \ge 0\}$. Here, $W = \{W_t, \ t \ge 0\}$ is a (standard) Wiener process, $M_t = \max_{0 \le s \le t} W_s$, and L_t is the local time of W at level zero and time t. In a recent paper [3], the author presents an elementary derivation of a discrete analogue of this result, for a symmetric simple random walk, which he then uses to derive Lévy's equivalence. The objective here is to point out that there is a trivariate version of Lévy's equivalence, which states that the processes $\{(M_{\downarrow} - W_{\downarrow}, M_{\downarrow}, \Theta_{\downarrow}), t \geq 0\}$ and $\{(|W_t|, L_t, T_t), t \ge 0\}$ have the same distributions, where $\Theta_t \in [0,t]$ is the time at which the maximum M, is attained, and T, is the last zero of W The proof depends on Tanaka's formula (cf. McKean (1969), page 68), in the time interval [0.1].

which says

$$L_{t} = |W_{t}| + \tilde{W}_{t} , t \ge 0,$$

where $\overline{W} = {\{W_t, t \ge 0\}}$ is a new Wiener process defined by

$$\tilde{w}_{t} = \int_{0}^{t} h(w_{s}) dw_{s} , t \ge 0,$$

with $h(\cdot) = -sign(\cdot)$ (cf. McKean (1969), page 29). Observe that

$$\bar{W}_{s} - \bar{W}_{t} = |W_{t}| - |W_{s}| - (L_{t} - L_{s}) \le |W_{t}|, s \in [0, t].$$

The inequality is an equality if and only if $s = T_t$. Thus


The inequality is an equality if and only if $s = T_t$. Thus

(1)
$$(\tilde{M}_t - \tilde{W}_t, \tilde{M}_t, \tilde{\Theta}_t) = (|W_t|, L_t, T_t)$$
, $t \ge 0$,

where $M_t = \max_{0 \le s \le t} W_s$, and $\Theta_t \in [0,t]$ is the time of the maximum. It should be emphasized that (1) is an almost sure identity in t for two trivariate stochastic processes. Consequently, $\{(M_t - W_t, M_t, \Theta_t), t \ge 0\}$ and $\{(|W_t|, L_t, T_t), t \ge 0\}$ have the same distributions as asserted.

REFERENCES

- [1] Knight, F. (1981), <u>Essentials of Browian Motion and Diffusion</u>, Mathematical Surveys No. 18 (American Math. Soc., Providence).
- [2] McKean, H.P. (1969), Stochastic Integrals, (Academic Press, New York).
- [3] Simons, G. (1983), A discrete analogue and elementary derivation of 'Lévy's equivalence' for Browian motion, <u>Statistics and Probability Letters</u> 1, 203-206.



END

FILMED

5-85

DTIC